

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

A metrological approach to evaluate epidemiological forecasting models

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1712371> since 2019-09-26T02:57:11Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

TOPIC: _Metrolog. traceability/Measurement uncertainty

Presentation preference: ☒ O ☐ P ☐ I ☐

A metrological approach to evaluate epidemiological forecasting models

Sanna Francesca¹⁾, Bellagarda Simone²⁾, Roggero Guido²⁾, Merlone Andrea²⁾

¹⁾ IMAMOTER-CNR - Istituto per le Macchine Agricole e Movimento Terra – Consiglio Nazionale Ricerche, Strada delle Cacce, 73, 10135, Turin, Italy – f.sanna@ima.to.cnr.it

²⁾ INRiM - Istituto Nazionale di Ricerca Metrologica, Strada delle Cacce, 91, 10135, Turin, Italy – s.bellagarda@inrim.it, g.roggero@inrim.it, a.merlone@inrim.it

Abstract: The evolve of the vineyard diseases, such as Grapevine Downy Mildew (*Plasmopara viticola*), are strictly depending by temperature, humidity and rain. The pathology is currently controlled with the massive use of fungicides, which has considerable economic costs, negative effects on environment, human health and wine quality.

For a correct defense against pathogen attacks, it is necessary to know the incubation period to act promptly and reduce the use of chemicals. In order to identify high-risk and fungicide sprays periods, several forecasting models have been proposed. These are useful tools which may assist in agricultural management risks and require accurate knowledge of meteorological variables such as temperature, humidity and precipitation. To date, the models used have improved the quality of the output data, but none of them considered the quality of the input data in terms of evaluation of measurement uncertainty and traceability of the sensors of the weather stations.

In situ calibration of weather stations installed in agricultural sites is usually performed by comparison. This procedure was metrologically evaluated and showed relevant weak points. Standard sensors are not always made to operate in open air, it is not possible to cover the whole range for the quantities, thus it is not possible to evaluate linearity and uncertainties for several sensors over the whole range and the evaluation of the mutual influences between parameters is not achievable. A calibration procedure for automatic weather stations for agrometeorological scope is proposed.

The aims of this study, part of the European MeteoMet project, are the improvement of the meteorological observations in field by disseminating the calibration methods, the implementation of traceability in agrometeorological measurements, and the improvement of the forecasting models by inclusion of traceable data and uncertainty components in the input values in order to reduce the use of chemicals in viticulture.